



ERA
Registry
Survival
Report
2023

ERA Registry

Amsterdam UMC, location AMC
Department of Medical Informatics
PO Box 22700
1100 DE Amsterdam
Netherlands

Telephone +31 20 566 7637

E-mail erareg@amsterdamumc.nl

Website www.era-online.org/research-education/era-registry

Further information (e.g. collaborating registries, methods) can be found in the ERA Registry annual report 2023, which is published on the ERA Registry website www.era-online.org/research-education/era-registry/annual-reports

ERA Registry
Survival Report 2023
Spain

Table of Contents

A5 SURVIVAL PROBABILITY

Table A.5.1	One-, two- and five-year survival probabilities by treatment modality and cohort	6
--------------------	--	---

B6 SURVIVAL PROBABILITY

Patient Survival on KRT (cohort 2014-2018 and cohort 2017-2021)

Table B.6.1	Incident KRT patients, from day 1, unadjusted	8
Table B.6.2	Incident KRT patients, from day 1, adjusted	10
Table B.6.3	Incident KRT patients, from day 91, unadjusted	12
Table B.6.4	Incident KRT patients, from day 91, adjusted	14

Patient Survival on Dialysis (cohort 2014-2018 and cohort 2017-2021)

Table B.6.5	Incident dialysis patients, from day 1, unadjusted	16
Table B.6.6	Incident dialysis patients, from day 1, adjusted	18
Table B.6.7	Incident dialysis patients, from day 91, unadjusted	20
Table B.6.8	Incident dialysis patients, from day 91, adjusted	22

Patient Survival after First Kidney Transplant (cohort 2014-2018 and cohort 2017-2021)

Table B.6.9	First transplant patients (deceased donor), from day of transplant, unadjusted	24
Table B.6.10	First transplant patients (deceased donor), from day of transplant, adjusted	26
Table B.6.11	First transplant patients (living donor), from day of transplant, unadjusted	28
Table B.6.12	First transplant patients (living donor), from day of transplant, adjusted	30

Graft Survival after First Kidney Transplant (cohort 2014-2018 and cohort 2017-2021)

Table B.6.13	First graft (deceased donor), from day of transplant, unadjusted	32
Table B.6.14	First graft (deceased donor), from day of transplant, adjusted	34
Table B.6.15	First graft (living donor), from day of transplant, unadjusted	36
Table B.6.16	First graft (living donor), from day of transplant, adjusted	38

Table A.5.1

One-, two- and five-year survival probabilities by treatment modality and cohort*from day 1 of KRT/ dialysis, or from day of transplant*

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
Patient survival on KRT					
Unadjusted	85.5 (85.3-85.7)	75.5 (75.3-75.7)	51.3 (51.1-51.6)	85.7 (85.5-85.9)	75.7 (75.5-75.9)
Adjusted †	88.3 (88.2-88.5)	79.4 (79.2-79.6)	54.0 (53.8-54.3)	88.3 (88.2-88.5)	79.3 (79.1-79.5)
Patient survival on dialysis					
Unadjusted	84.4 (84.3-84.6)	72.8 (72.6-73.0)	41.4 (41.1-41.6)	84.7 (84.6-84.9)	73.3 (73.1-73.5)
Adjusted †	86.7 (86.6-86.9)	76.4 (76.2-76.6)	46.7 (46.4-47.0)	87.1 (87.0-87.3)	77.0 (76.8-77.2)
Patient survival after first kidney transplantation (deceased donor)					
Unadjusted	96.3 (96.1-96.5)	94.1 (93.8-94.3)	84.3 (84.0-84.7)	95.8 (95.6-96.0)	93.0 (92.8-93.3)
Adjusted ††	98.2 (98.1-98.3)	97.0 (96.9-97.2)	91.6 (91.3-91.9)	98.1 (98.0-98.2)	96.7 (96.5-96.9)
Graft survival after first kidney transplantation (deceased donor)					
Unadjusted	91.1 (90.8-91.4)	87.7 (87.4-88.0)	75.4 (75.0-75.8)	90.9 (90.6-91.2)	87.2 (86.9-87.6)
Adjusted ††	93.3 (93.0-93.5)	90.6 (90.4-90.9)	80.6 (80.2-81.0)	93.4 (93.1-93.6)	90.6 (90.3-90.9)
Patient survival after first kidney transplantation (living donor)					
Unadjusted	98.8 (98.6-99.0)	98.0 (97.8-98.2)	94.0 (93.6-94.4)	98.8 (98.6-99.0)	97.9 (97.6-98.1)
Adjusted ††	99.1 (99.0-99.3)	98.5 (98.3-98.7)	95.2 (94.8-95.6)	99.1 (98.9-99.3)	98.4 (98.1-98.6)
Graft survival after first kidney transplantation (living donor)					
Unadjusted	96.5 (96.2-96.8)	95.0 (94.6-95.3)	88.0 (87.4-88.5)	96.9 (96.6-97.2)	95.3 (94.9-95.7)
Adjusted ††	96.4 (96.1-96.8)	94.8 (94.4-95.2)	87.6 (87.0-88.2)	96.8 (96.5-97.2)	95.1 (94.7-95.5)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

† Analyses were adjusted using fixed values: age (67 years), sex (63% male) and PRD (24% diabetes mellitus, 19% hypertension/renal vascular disease, 11% glomerulonephritis and 46% other causes)

†† Analyses were adjusted using fixed values: age (50 years), sex (63% male) and PRD (14% diabetes mellitus, 10% hypertension/renal vascular disease, 23% glomerulonephritis and 53% other causes)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
Patient survival on KRT					
Unadjusted	89.7 (89.4-90.0)	81.1 (80.6-81.5)	58.1 (57.6-58.7)	89.6 (89.3-90.0)	81.1 (80.7-81.5)
Adjusted †	91.5 (91.2-91.8)	83.9 (83.5-84.3)	61.0 (60.4-61.6)	91.3 (91.1-91.6)	83.7 (83.3-84.1)
Patient survival on dialysis					
Unadjusted	89.0 (88.7-89.4)	78.8 (78.3-79.3)	46.5 (45.8-47.2)	89.0 (88.6-89.3)	78.9 (78.5-79.4)
Adjusted †	90.3 (90.0-90.7)	81.4 (80.9-81.9)	52.7 (52.0-53.4)	90.5 (90.2-90.8)	81.8 (81.3-82.3)
Patient survival after first kidney transplantation (deceased donor)					
Unadjusted	95.4 (95.0-95.8)	93.2 (92.8-93.7)	82.7 (82.0-83.4)	95.2 (94.8-95.6)	92.5 (91.9-92.9)
Adjusted ††	98.1 (97.9-98.3)	97.2 (97.0-97.5)	92.2 (91.7-92.8)	98.1 (97.9-98.3)	97.0 (96.7-97.3)
Graft survival after first kidney transplantation (deceased donor)					
Unadjusted	88.6 (88.0-89.2)	85.3 (84.7-86.0)	72.5 (71.7-73.3)	88.7 (88.1-89.3)	85.0 (84.3-85.7)
Adjusted ††	92.0 (91.5-92.5)	89.6 (89.0-90.1)	79.6 (78.7-80.4)	92.1 (91.6-92.7)	89.5 (88.8-90.1)
Patient survival after first kidney transplantation (living donor)					
Unadjusted	99.3 (98.7-99.6)	98.5 (97.7-99.0)	95.7 (94.6-96.7)	99.2 (98.5-99.6)	98.3 (97.4-98.9)
Adjusted ††	99.5 (99.2-99.8)	98.9 (98.4-99.4)	97.0 (96.1-97.9)	99.6 (99.3-99.9)	99.2 (98.7-99.7)
Graft survival after first kidney transplantation (living donor)					
Unadjusted	95.7 (94.5-96.6)	94.1 (92.7-95.2)	89.1 (87.3-90.6)	96.2 (95.0-97.2)	94.6 (93.2-95.8)
Adjusted ††	95.5 (94.4-96.6)	93.8 (92.5-95.1)	88.6 (86.9-90.3)	96.3 (95.2-97.4)	94.7 (93.5-96.0)

† Analyses were adjusted using fixed values: age (67 years), sex (63% male) and PRD (24% diabetes mellitus, 19% hypertension/renal vascular disease, 11% glomerulonephritis and 46% other causes)

†† Analyses were adjusted using fixed values: age (50 years), sex (63% male) and PRD (14% diabetes mellitus, 10% hypertension/renal vascular disease, 23% glomerulonephritis and 53% other causes)

Table B.6.1
Incident KRT patients, from day 1, unadjusted
by age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)						
	cohort 2014 - 2018				cohort 2017 - 2021		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.5 (99.1-99.7)	98.0 (97.4-98.5)	97.2 (96.4-97.8)	95.0 (94.0-95.8)	99.3 (98.8-99.5)	97.8 (97.1-98.3)	97.1 (96.4-97.7)
20-44	99.2 (99.0-99.3)	97.0 (96.8-97.3)	94.8 (94.5-95.1)	89.0 (88.5-89.5)	99.1 (99.0-99.2)	96.8 (96.6-97.1)	94.3 (93.9-94.6)
45-64	97.5 (97.4-97.7)	92.0 (91.8-92.2)	85.8 (85.5-86.1)	69.1 (68.7-69.5)	97.6 (97.5-97.8)	92.2 (92.0-92.4)	85.9 (85.6-86.2)
65-74	95.2 (95.0-95.4)	84.9 (84.6-85.2)	74.1 (73.7-74.5)	46.1 (45.6-46.5)	95.1 (94.9-95.3)	85.0 (84.7-85.3)	74.2 (73.8-74.6)
75+	91.5 (91.3-91.7)	75.8 (75.4-76.1)	60.0 (59.6-60.4)	25.4 (25.0-25.8)	91.6 (91.3-91.8)	76.4 (76.0-76.7)	60.8 (60.4-61.2)
Male	95.2 (95.1-95.3)	85.3 (85.1-85.5)	75.0 (74.7-75.2)	50.1 (49.8-50.3)	95.1 (95.0-95.3)	85.4 (85.2-85.6)	75.0 (74.8-75.3)
Female	95.2 (95.0-95.3)	85.9 (85.6-86.2)	76.5 (76.1-76.8)	53.7 (53.3-54.1)	95.2 (95.0-95.4)	86.3 (86.0-86.6)	76.9 (76.5-77.2)
Diabetes	96.1 (95.9-96.3)	86.5 (86.2-86.8)	74.8 (74.3-75.2)	44.5 (44.0-45.0)	96.0 (95.8-96.2)	86.4 (86.1-86.7)	74.5 (74.1-74.9)
Hypertension / renal vascular disease	95.0 (94.8-95.2)	84.4 (84.0-84.7)	72.6 (72.1-73.1)	43.4 (42.9-44.0)	95.1 (94.9-95.3)	84.8 (84.4-85.2)	73.0 (72.5-73.5)
Glomerulonephritis	97.9 (97.7-98.0)	93.2 (92.8-93.5)	88.1 (87.6-88.5)	72.6 (72.0-73.3)	98.1 (97.9-98.2)	93.7 (93.3-94.0)	88.4 (88.0-88.9)
Other causes	94.4 (94.2-94.5)	83.9 (83.7-84.2)	74.3 (74.0-74.6)	53.1 (52.8-53.5)	94.4 (94.2-94.6)	84.3 (84.0-84.5)	74.8 (74.5-75.1)
All	95.2 (95.1-95.3)	85.5 (85.3-85.7)	75.5 (75.3-75.7)	51.3 (51.1-51.6)	95.2 (95.1-95.3)	85.7 (85.5-85.9)	75.7 (75.5-75.9)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)						
	cohort 2014 - 2018				cohort 2017 - 2021		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.4 (97.6-99.9)	98.2 (96.0-99.2)	97.9 (95.6-99.0)	96.7 (94.1-98.1)	99.2 (97.5-99.7)	98.3 (96.4-99.3)	98.3 (96.4-99.3)
20-44	99.4 (99.1-99.6)	97.9 (97.4-98.4)	96.3 (95.6-96.9)	92.3 (91.3-93.2)	99.1 (98.8-99.4)	97.7 (97.1-98.2)	96.1 (95.4-96.7)
45-64	98.4 (98.1-98.6)	94.1 (93.6-94.6)	89.0 (88.3-89.6)	74.0 (73.1-74.9)	98.6 (98.4-98.8)	94.4 (93.9-94.8)	89.5 (88.9-90.1)
65-74	96.8 (96.4-97.1)	88.4 (87.7-89.0)	78.6 (77.7-79.5)	52.1 (51.0-53.1)	96.7 (96.3-97.1)	87.9 (87.2-88.5)	78.4 (77.5-79.2)
75+	95.1 (94.6-95.5)	83.1 (82.4-83.9)	69.2 (68.3-70.2)	34.0 (33.0-35.0)	95.5 (95.1-95.9)	83.6 (82.9-84.3)	69.7 (68.8-70.6)
Male	96.9 (96.7-97.2)	89.3 (88.9-89.7)	80.3 (79.7-80.8)	56.1 (55.4-56.7)	97.1 (96.9-97.3)	89.1 (88.6-89.5)	80.1 (79.6-80.6)
Female	97.4 (97.0-97.7)	90.5 (89.9-91.1)	82.8 (82.0-83.5)	62.3 (61.4-63.3)	97.4 (97.1-97.7)	90.8 (90.2-91.3)	83.1 (82.4-83.8)
Diabetes	97.4 (97.0-97.7)	89.3 (88.6-90.0)	78.7 (77.8-79.6)	50.0 (48.9-51.1)	97.6 (97.2-97.9)	88.9 (88.3-89.6)	78.8 (77.9-79.6)
Hypertension / renal vascular disease	97.4 (96.8-97.8)	89.9 (88.9-90.8)	79.9 (78.7-81.1)	52.3 (50.7-53.8)	97.2 (96.6-97.6)	89.2 (88.2-90.1)	78.9 (77.6-80.1)
Glomerulonephritis	98.5 (98.0-98.8)	94.2 (93.4-94.9)	89.4 (88.4-90.4)	75.0 (73.7-76.4)	98.7 (98.4-99.0)	95.1 (94.4-95.7)	90.2 (89.3-91.1)
Other causes	96.5 (96.2-96.8)	88.8 (88.3-89.3)	80.6 (80.0-81.2)	59.8 (59.0-60.6)	96.8 (96.5-97.1)	89.0 (88.5-89.5)	80.9 (80.3-81.5)
All	97.1 (96.9-97.3)	89.7 (89.4-90.0)	81.1 (80.6-81.5)	58.1 (57.6-58.7)	97.2 (97.0-97.4)	89.6 (89.3-90.0)	81.1 (80.7-81.5)

Table B.6.2
Incident KRT patients, from day 1, adjusted
adjusted for age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)						
	cohort 2014 - 2018				cohort 2017 - 2021		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.5 (99.2-99.8)	97.9 (97.3-98.5)	97.0 (96.3-97.7)	94.7 (93.7-95.6)	99.2 (98.9-99.6)	97.7 (97.0-98.3)	97.0 (96.3-97.7)
20-44	99.2 (99.1-99.3)	97.0 (96.7-97.2)	94.7 (94.3-95.0)	88.6 (88.1-89.1)	99.1 (99.0-99.3)	96.8 (96.5-97.1)	94.2 (93.8-94.5)
45-64	97.7 (97.6-97.8)	92.3 (92.1-92.5)	86.2 (85.9-86.5)	69.4 (69.0-69.8)	97.8 (97.7-97.9)	92.5 (92.3-92.8)	86.3 (86.0-86.6)
65-74	95.5 (95.4-95.7)	85.7 (85.4-86.0)	75.2 (74.8-75.6)	47.4 (46.9-47.9)	95.4 (95.2-95.6)	85.7 (85.4-86.0)	75.2 (74.9-75.6)
75+	91.8 (91.6-92.1)	76.5 (76.1-76.8)	60.9 (60.5-61.3)	26.2 (25.8-26.6)	92.0 (91.8-92.2)	77.3 (76.9-77.6)	61.9 (61.5-62.3)
Male	96.3 (96.2-96.4)	88.1 (87.9-88.3)	78.8 (78.6-79.0)	52.5 (52.2-52.8)	96.2 (96.1-96.3)	88.0 (87.8-88.2)	78.6 (78.4-78.9)
Female	96.3 (96.1-96.4)	88.6 (88.4-88.9)	80.2 (79.9-80.5)	56.5 (56.1-56.9)	96.3 (96.1-96.4)	88.8 (88.5-89.0)	80.2 (79.9-80.5)
Diabetes	96.7 (96.5-96.8)	88.4 (88.1-88.7)	77.6 (77.3-78.0)	46.6 (46.1-47.1)	96.6 (96.4-96.8)	88.3 (88.0-88.5)	77.4 (77.0-77.8)
Hypertension / renal vascular disease	96.9 (96.8-97.1)	90.0 (89.7-90.3)	81.6 (81.3-82.0)	55.9 (55.4-56.4)	96.8 (96.7-97.0)	89.7 (89.4-90.0)	81.0 (80.6-81.4)
Glomerulonephritis	97.6 (97.3-97.8)	92.0 (91.6-92.4)	85.8 (85.3-86.3)	65.6 (64.9-66.3)	97.8 (97.6-98.0)	92.6 (92.3-93.0)	86.3 (85.8-86.8)
Other causes	95.3 (95.2-95.5)	86.2 (86.0-86.4)	77.1 (76.8-77.4)	53.8 (53.5-54.2)	95.3 (95.2-95.4)	86.3 (86.1-86.5)	77.3 (77.0-77.6)
All	96.3 (96.2-96.4)	88.3 (88.2-88.5)	79.4 (79.2-79.6)	54.0 (53.8-54.3)	96.2 (96.1-96.3)	88.3 (88.2-88.5)	79.3 (79.1-79.5)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)						
	cohort 2014 - 2018				cohort 2017 - 2021		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.4 (98.5-100)	98.1 (96.6-99.6)	97.8 (96.1-99.4)	96.4 (94.4-98.5)	99.1 (98.2-100)	98.3 (96.9-99.7)	98.3 (96.9-99.7)
20-44	99.4 (99.1-99.7)	97.8 (97.3-98.4)	96.2 (95.5-96.9)	91.9 (91.0-92.9)	99.2 (98.9-99.5)	97.7 (97.1-98.2)	96.0 (95.3-96.7)
45-64	98.4 (98.2-98.7)	94.2 (93.8-94.7)	89.2 (88.6-89.8)	74.3 (73.4-75.1)	98.7 (98.5-98.9)	94.5 (94.1-95.0)	89.7 (89.1-90.3)
65-74	97.0 (96.6-97.3)	88.9 (88.3-89.6)	79.6 (78.7-80.4)	53.6 (52.5-54.7)	96.9 (96.6-97.3)	88.4 (87.8-89.1)	79.1 (78.3-80.0)
75+	95.2 (94.8-95.6)	83.6 (82.8-84.3)	69.9 (69.0-70.9)	34.9 (33.9-35.9)	95.7 (95.3-96.1)	84.3 (83.6-85.0)	70.7 (69.8-71.6)
Male	97.5 (97.3-97.7)	91.1 (90.7-91.4)	83.0 (82.6-83.5)	58.6 (57.9-59.3)	97.6 (97.5-97.8)	90.8 (90.4-91.2)	82.7 (82.3-83.2)
Female	97.8 (97.6-98.1)	92.1 (91.6-92.6)	85.3 (84.7-86.0)	65.0 (64.1-66.0)	97.9 (97.6-98.1)	92.2 (91.8-92.7)	85.3 (84.7-86.0)
Diabetes	97.8 (97.5-98.1)	91.1 (90.5-91.6)	81.7 (80.9-82.6)	53.7 (52.5-54.8)	98.0 (97.7-98.2)	90.7 (90.1-91.3)	81.8 (81.0-82.6)
Hypertension / renal vascular disease	98.3 (98.0-98.6)	93.4 (92.8-94.0)	86.4 (85.5-87.3)	63.5 (62.1-64.9)	98.1 (97.7-98.4)	92.5 (91.9-93.2)	84.9 (83.9-85.9)
Glomerulonephritis	98.2 (97.8-98.7)	93.2 (92.4-94.1)	87.5 (86.4-88.7)	69.1 (67.4-70.7)	98.6 (98.2-99.0)	94.4 (93.7-95.2)	88.7 (87.7-89.8)
Other causes	97.1 (96.9-97.3)	90.4 (89.9-90.8)	82.9 (82.3-83.5)	61.3 (60.5-62.1)	97.2 (97.0-97.5)	90.4 (89.9-90.8)	82.8 (82.2-83.4)
All	97.6 (97.5-97.8)	91.5 (91.2-91.8)	83.9 (83.5-84.3)	61.0 (60.4-61.6)	97.7 (97.6-97.9)	91.3 (91.1-91.6)	83.7 (83.3-84.1)

Table B.6.3
Incident KRT patients, from day 91, unadjusted
by age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	98.3 (97.7-98.8)	97.6 (96.9-98.1)	95.4 (94.4-96.1)	98.2 (97.6-98.7)	97.7 (97.0-98.3)
20-44	97.2 (97.0-97.4)	95.0 (94.7-95.3)	89.4 (88.9-89.8)	97.1 (96.8-97.3)	94.6 (94.2-94.9)
45-64	92.8 (92.6-93.1)	86.6 (86.3-86.9)	69.7 (69.3-70.1)	92.8 (92.6-93.0)	86.6 (86.3-86.9)
65-74	86.5 (86.2-86.8)	75.1 (74.7-75.5)	46.6 (46.1-47.0)	86.7 (86.4-87.0)	75.6 (75.2-76.0)
75+	78.7 (78.4-79.1)	61.8 (61.4-62.3)	25.8 (25.4-26.2)	79.4 (79.1-79.8)	62.7 (62.3-63.1)
Male	87.0 (86.8-87.2)	76.3 (76.0-76.5)	51.1 (50.8-51.4)	87.1 (86.9-87.3)	76.5 (76.2-76.7)
Female	87.9 (87.7-88.2)	78.2 (77.9-78.6)	55.0 (54.6-55.4)	88.3 (88.1-88.6)	78.7 (78.4-79.0)
Diabetes	87.1 (86.8-87.5)	74.9 (74.4-75.3)	44.4 (43.9-44.9)	87.2 (86.9-87.6)	74.9 (74.4-75.3)
Hypertension / renal vascular disease	85.9 (85.5-86.2)	73.7 (73.2-74.2)	43.9 (43.3-44.5)	86.1 (85.7-86.5)	74.2 (73.7-74.7)
Glomerulonephritis	94.0 (93.6-94.3)	88.7 (88.2-89.1)	73.1 (72.5-73.8)	94.1 (93.7-94.4)	88.9 (88.5-89.3)
Other causes	86.5 (86.3-86.8)	76.6 (76.3-76.9)	55.1 (54.8-55.5)	86.8 (86.6-87.1)	77.1 (76.8-77.4)
All	87.4 (87.2-87.5)	77.0 (76.8-77.2)	52.5 (52.2-52.7)	87.5 (87.4-87.7)	77.3 (77.1-77.5)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	98.8 (96.8-99.5)	98.5 (96.4-99.4)	97.2 (94.8-98.6)	99.2 (97.4-99.7)	99.2 (97.4-99.7)
20-44	97.9 (97.4-98.4)	96.7 (96.0-97.3)	92.5 (91.5-93.4)	98.1 (97.5-98.5)	96.8 (96.1-97.3)
45-64	94.4 (93.9-94.8)	89.3 (88.6-89.9)	74.1 (73.2-75.0)	94.4 (94.0-94.9)	89.6 (89.0-90.2)
65-74	89.3 (88.6-89.9)	78.8 (77.9-79.7)	51.9 (50.8-53.0)	88.5 (87.8-89.1)	78.6 (77.8-79.5)
75+	83.7 (82.9-84.5)	69.2 (68.2-70.1)	33.1 (32.1-34.1)	84.1 (83.3-84.8)	69.9 (68.9-70.8)
Male	90.0 (89.6-90.4)	80.4 (79.8-80.9)	56.1 (55.4-56.7)	89.5 (89.1-89.9)	80.4 (79.9-80.9)
Female	90.9 (90.3-91.4)	83.4 (82.7-84.1)	62.6 (61.6-63.5)	91.2 (90.6-91.7)	83.5 (82.8-84.2)
Diabetes	89.1 (88.3-89.8)	78.4 (77.4-79.3)	49.2 (48.0-50.3)	88.8 (88.1-89.4)	78.1 (77.2-79.0)
Hypertension / renal vascular disease	90.0 (89.0-90.9)	79.6 (78.3-80.8)	51.4 (49.8-53.0)	89.5 (88.5-90.4)	79.1 (77.8-80.3)
Glomerulonephritis	94.6 (93.8-95.3)	89.6 (88.6-90.5)	75.1 (73.7-76.5)	95.2 (94.5-95.8)	90.1 (89.1-91.0)
Other causes	89.9 (89.4-90.4)	81.5 (80.8-82.1)	60.6 (59.8-61.4)	89.8 (89.3-90.2)	81.8 (81.1-82.4)
All	90.3 (89.9-90.6)	81.4 (81.0-81.8)	58.2 (57.6-58.8)	90.1 (89.7-90.4)	81.4 (81.0-81.8)

Table B.6.4
Incident KRT patients, from day 91, adjusted
adjusted for age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	98.2 (97.7-98.8)	97.4 (96.8-98.1)	95.0 (94.1-96.0)	98.1 (97.5-98.7)	97.6 (96.9-98.2)
20-44	97.1 (96.8-97.3)	94.8 (94.5-95.2)	88.8 (88.4-89.3)	97.0 (96.7-97.2)	94.4 (94.1-94.8)
45-64	93.0 (92.8-93.3)	86.9 (86.6-87.2)	69.9 (69.5-70.3)	93.0 (92.8-93.2)	86.9 (86.6-87.2)
65-74	87.2 (86.8-87.5)	76.2 (75.8-76.6)	47.9 (47.4-48.4)	87.3 (87.0-87.6)	76.5 (76.2-76.9)
75+	79.3 (79.0-79.7)	62.7 (62.3-63.1)	26.6 (26.2-27.0)	80.1 (79.7-80.4)	63.7 (63.3-64.1)
Male	89.3 (89.1-89.5)	79.6 (79.4-79.9)	52.9 (52.5-53.2)	89.2 (89.1-89.4)	79.6 (79.3-79.8)
Female	90.1 (89.9-90.3)	81.4 (81.1-81.7)	57.1 (56.7-57.5)	90.2 (89.9-90.4)	81.4 (81.1-81.7)
Diabetes	88.8 (88.6-89.1)	77.5 (77.1-77.9)	46.1 (45.6-46.6)	88.9 (88.6-89.2)	77.5 (77.1-77.9)
Hypertension / renal vascular disease	90.9 (90.7-91.2)	82.3 (81.9-82.6)	55.9 (55.3-56.4)	90.6 (90.3-90.8)	81.6 (81.2-82.0)
Glomerulonephritis	92.9 (92.5-93.3)	86.3 (85.8-86.8)	65.7 (65.0-66.5)	93.0 (92.6-93.4)	86.7 (86.2-87.2)
Other causes	88.2 (88.0-88.4)	78.8 (78.5-79.0)	55.0 (54.6-55.4)	88.4 (88.1-88.6)	79.1 (78.8-79.4)
All	89.6 (89.5-89.8)	80.3 (80.1-80.5)	54.5 (54.2-54.7)	89.6 (89.5-89.7)	80.3 (80.1-80.5)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	98.7 (97.4-100)	98.4 (96.9-99.8)	97.0 (95.1-99.0)	99.1 (98.1-100)	99.1 (98.1-100)
20-44	97.8 (97.3-98.3)	96.5 (95.9-97.2)	92.1 (91.2-93.1)	98.0 (97.5-98.5)	96.6 (95.9-97.3)
45-64	94.5 (94.0-94.9)	89.5 (88.8-90.1)	74.4 (73.5-75.3)	94.5 (94.1-95.0)	89.8 (89.2-90.4)
65-74	89.8 (89.1-90.4)	79.8 (78.9-80.6)	53.5 (52.4-54.5)	88.9 (88.3-89.6)	79.4 (78.5-80.2)
75+	84.1 (83.4-84.9)	69.9 (68.9-70.8)	34.0 (33.0-35.1)	84.7 (83.9-85.4)	70.8 (69.9-71.7)
Male	91.6 (91.2-92.0)	83.1 (82.6-83.6)	58.2 (57.5-58.9)	91.1 (90.8-91.5)	83.0 (82.5-83.5)
Female	92.4 (91.9-92.9)	85.8 (85.1-86.4)	65.0 (64.1-66.0)	92.5 (92.1-93.0)	85.6 (85.0-86.3)
Diabetes	90.8 (90.2-91.4)	81.4 (80.6-82.2)	52.6 (51.5-53.8)	90.6 (90.0-91.2)	81.3 (80.5-82.1)
Hypertension / renal vascular disease	93.5 (92.8-94.1)	86.2 (85.3-87.1)	62.6 (61.2-64.0)	92.8 (92.1-93.5)	85.2 (84.2-86.2)
Glomerulonephritis	93.7 (92.8-94.5)	87.6 (86.5-88.8)	68.9 (67.2-70.5)	94.5 (93.7-95.2)	88.5 (87.4-89.6)
Other causes	91.3 (90.9-91.7)	83.5 (82.9-84.1)	61.7 (60.9-62.5)	91.0 (90.6-91.5)	83.6 (83.0-84.2)
All	91.9 (91.6-92.2)	84.1 (83.7-84.5)	60.7 (60.1-61.3)	91.7 (91.4-92.0)	84.0 (83.6-84.4)

Table B.6.5
Incident dialysis patients, from day 1, unadjusted
by age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)						
	cohort 2014 - 2018				cohort 2017 - 2021		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.4 (98.9-99.7)	97.3 (96.3-98.0)	95.5 (94.1-96.6)	87.1 (83.1-90.2)	99.3 (98.8-99.6)	97.2 (96.3-97.9)	96.2 (95.0-97.1)
20-44	99.1 (98.9-99.2)	96.4 (96.1-96.7)	92.8 (92.3-93.2)	79.6 (78.6-80.5)	99.0 (98.8-99.2)	96.2 (95.8-96.5)	92.3 (91.8-92.7)
45-64	97.4 (97.2-97.5)	91.2 (90.9-91.4)	83.6 (83.2-83.9)	58.9 (58.4-59.4)	97.5 (97.4-97.6)	91.4 (91.2-91.7)	83.9 (83.5-84.2)
65-74	95.1 (94.9-95.3)	84.4 (84.0-84.7)	72.8 (72.3-73.2)	41.0 (40.5-41.5)	94.9 (94.7-95.1)	84.4 (84.1-84.7)	72.9 (72.5-73.4)
75+	91.5 (91.2-91.7)	75.7 (75.3-76.0)	59.7 (59.3-60.1)	24.7 (24.3-25.0)	91.5 (91.3-91.7)	76.2 (75.9-76.6)	60.5 (60.1-60.9)
Male	94.9 (94.8-95.1)	84.2 (84.0-84.5)	72.3 (72.0-72.6)	39.9 (39.6-40.3)	94.9 (94.8-95.0)	84.5 (84.2-84.7)	72.7 (72.4-72.9)
Female	94.9 (94.7-95.0)	84.8 (84.5-85.1)	73.8 (73.4-74.2)	44.0 (43.5-44.4)	94.9 (94.7-95.1)	85.3 (85.0-85.5)	74.4 (74.1-74.8)
Diabetes	96.0 (95.8-96.2)	86.1 (85.8-86.4)	73.5 (73.1-73.9)	38.6 (38.1-39.2)	95.9 (95.7-96.1)	86.1 (85.7-86.4)	73.5 (73.1-73.9)
Hypertension / renal vascular disease	94.9 (94.7-95.2)	83.9 (83.5-84.3)	71.3 (70.8-71.9)	37.8 (37.3-38.4)	95.0 (94.8-95.3)	84.4 (84.0-84.8)	71.8 (71.3-72.4)
Glomerulonephritis	97.6 (97.4-97.9)	92.2 (91.8-92.6)	85.4 (84.9-86.0)	59.6 (58.7-60.5)	97.9 (97.7-98.1)	92.8 (92.5-93.2)	86.0 (85.5-86.5)
Other causes	93.9 (93.8-94.1)	82.3 (82.0-82.6)	70.7 (70.3-71.0)	41.6 (41.2-42.0)	94.0 (93.8-94.2)	82.8 (82.5-83.1)	71.6 (71.2-71.9)
All	94.9 (94.8-95.0)	84.4 (84.3-84.6)	72.8 (72.6-73.0)	41.4 (41.1-41.6)	94.9 (94.8-95.0)	84.7 (84.6-84.9)	73.3 (73.1-73.5)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)						
	cohort 2014 - 2018				cohort 2017 - 2021		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.5 (96.8-99.9)	97.4 (93.9-98.9)	95.9 (90.1-98.3)	92.8 (81.6-97.3)	99.2 (96.8-99.8)	97.5 (94.0-99.0)	97.5 (94.0-99.0)
20-44	99.3 (99.0-99.6)	97.6 (96.9-98.1)	95.1 (94.1-96.0)	84.6 (82.1-86.7)	99.1 (98.7-99.4)	97.4 (96.7-97.9)	95.0 (94.0-95.9)
45-64	98.4 (98.1-98.6)	93.7 (93.1-94.2)	87.1 (86.3-87.9)	61.5 (60.0-62.9)	98.6 (98.4-98.8)	93.9 (93.4-94.4)	87.8 (87.0-88.5)
65-74	96.7 (96.3-97.1)	88.0 (87.3-88.7)	77.3 (76.4-78.3)	45.4 (44.1-46.6)	96.7 (96.3-97.0)	87.5 (86.7-88.1)	77.1 (76.2-78.0)
75+	95.1 (94.6-95.5)	83.1 (82.3-83.8)	68.9 (68.0-69.9)	32.9 (31.9-33.9)	95.5 (95.0-95.9)	83.5 (82.7-84.2)	69.4 (68.4-70.3)
Male	96.8 (96.6-97.1)	88.6 (88.2-89.1)	77.9 (77.3-78.5)	44.2 (43.3-45.0)	97.0 (96.8-97.2)	88.4 (88.0-88.9)	77.9 (77.3-78.5)
Female	97.2 (96.9-97.5)	89.9 (89.2-90.4)	80.7 (79.8-81.5)	51.3 (50.0-52.5)	97.3 (96.9-97.6)	90.1 (89.5-90.7)	81.1 (80.2-81.9)
Diabetes	97.3 (97.0-97.7)	89.1 (88.3-89.8)	77.7 (76.7-78.6)	42.7 (41.4-44.0)	97.5 (97.2-97.9)	88.7 (88.0-89.3)	77.8 (76.8-78.7)
Hypertension / renal vascular disease	97.3 (96.8-97.8)	89.5 (88.5-90.5)	78.6 (77.3-79.9)	44.4 (42.6-46.1)	97.1 (96.5-97.6)	88.8 (87.8-89.8)	77.4 (76.0-78.7)
Glomerulonephritis	98.3 (97.9-98.7)	93.6 (92.7-94.4)	87.1 (85.9-88.3)	61.4 (59.0-63.6)	98.7 (98.2-99.0)	94.6 (93.8-95.3)	88.3 (87.1-89.4)
Other causes	96.4 (96.0-96.7)	87.9 (87.3-88.4)	77.8 (77.1-78.5)	47.1 (46.0-48.1)	96.7 (96.4-96.9)	88.2 (87.6-88.7)	78.2 (77.5-78.9)
All	97.0 (96.8-97.2)	89.0 (88.7-89.4)	78.8 (78.3-79.3)	46.5 (45.8-47.2)	97.1 (96.9-97.3)	89.0 (88.6-89.3)	78.9 (78.5-79.4)

Table B.6.6
Incident dialysis patients, from day 1, adjusted
adjusted for age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)						
	cohort 2014 - 2018				cohort 2017 - 2021		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.4 (99.0-99.8)	97.2 (96.3-98.0)	95.3 (94.1-96.6)	86.7 (83.2-90.4)	99.3 (98.9-99.7)	97.1 (96.2-98.0)	96.0 (94.9-97.1)
20-44	99.1 (99.0-99.3)	96.4 (96.1-96.7)	92.7 (92.2-93.2)	79.3 (78.4-80.3)	99.0 (98.9-99.2)	96.2 (95.8-96.5)	92.2 (91.8-92.7)
45-64	97.5 (97.4-97.7)	91.5 (91.2-91.7)	84.0 (83.6-84.3)	59.5 (59.0-60.1)	97.7 (97.6-97.8)	91.8 (91.6-92.1)	84.4 (84.0-84.7)
65-74	95.4 (95.2-95.6)	85.0 (84.7-85.4)	73.6 (73.2-74.1)	42.1 (41.7-42.7)	95.3 (95.1-95.5)	85.1 (84.8-85.4)	73.9 (73.5-74.3)
75+	91.8 (91.5-92.0)	76.2 (75.9-76.6)	60.4 (60.0-60.8)	25.2 (24.8-25.6)	91.9 (91.7-92.1)	77.0 (76.7-77.4)	61.4 (61.0-61.8)
Male	95.8 (95.7-95.9)	86.5 (86.3-86.7)	75.8 (75.5-76.1)	45.1 (44.7-45.4)	95.8 (95.7-95.9)	86.8 (86.6-87.0)	76.4 (76.1-76.6)
Female	95.7 (95.6-95.9)	87.0 (86.8-87.3)	77.3 (77.0-77.7)	49.5 (49.0-49.9)	95.9 (95.7-96.0)	87.6 (87.4-87.9)	78.1 (77.7-78.4)
Diabetes	96.4 (96.2-96.6)	87.4 (87.1-87.7)	75.7 (75.3-76.1)	41.6 (41.1-42.1)	96.4 (96.2-96.6)	87.5 (87.2-87.8)	76.0 (75.6-76.4)
Hypertension / renal vascular disease	96.4 (96.2-96.6)	88.4 (88.1-88.7)	78.6 (78.2-79.1)	49.0 (48.4-49.6)	96.5 (96.3-96.6)	88.6 (88.3-88.9)	78.8 (78.4-79.2)
Glomerulonephritis	97.3 (97.1-97.6)	91.2 (90.8-91.7)	84.0 (83.4-84.6)	57.5 (56.5-58.4)	97.6 (97.4-97.9)	92.0 (91.6-92.4)	84.6 (84.0-85.2)
Other causes	94.7 (94.5-94.8)	84.3 (84.0-84.5)	73.7 (73.4-74.0)	45.9 (45.5-46.3)	94.8 (94.7-95.0)	84.9 (84.6-85.1)	74.7 (74.4-75.0)
All	95.8 (95.7-95.9)	86.7 (86.6-86.9)	76.4 (76.2-76.6)	46.7 (46.4-47.0)	95.9 (95.8-96.0)	87.1 (87.0-87.3)	77.0 (76.8-77.2)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)						
	cohort 2014 - 2018				cohort 2017 - 2021		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.5 (98.6-100)	97.3 (95.0-99.7)	95.7 (92.0-99.6)	92.6 (85.7-100)	99.2 (98.0-100)	97.4 (95.2-99.7)	97.4 (95.2-99.7)
20-44	99.3 (99.0-99.6)	97.5 (96.9-98.2)	95.0 (94.1-96.0)	84.4 (82.2-86.8)	99.1 (98.8-99.5)	97.3 (96.7-98.0)	95.0 (94.0-95.9)
45-64	98.4 (98.2-98.7)	93.8 (93.3-94.3)	87.5 (86.7-88.2)	62.3 (60.9-63.7)	98.7 (98.5-98.9)	94.1 (93.6-94.6)	88.2 (87.5-88.9)
65-74	96.9 (96.5-97.3)	88.5 (87.8-89.2)	78.2 (77.3-79.1)	46.9 (45.6-48.2)	96.9 (96.5-97.2)	88.0 (87.3-88.6)	77.9 (77.0-78.8)
75+	95.1 (94.7-95.6)	83.4 (82.6-84.2)	69.5 (68.6-70.5)	33.6 (32.6-34.6)	95.7 (95.3-96.1)	84.1 (83.4-84.8)	70.2 (69.3-71.2)
Male	97.2 (97.0-97.4)	89.9 (89.5-90.3)	80.3 (79.8-80.9)	49.9 (49.0-50.8)	97.5 (97.3-97.7)	89.9 (89.5-90.3)	80.7 (80.2-81.3)
Female	97.5 (97.3-97.8)	91.0 (90.5-91.6)	83.0 (82.3-83.8)	57.5 (56.4-58.7)	97.7 (97.4-97.9)	91.4 (90.9-92.0)	83.6 (82.8-84.3)
Diabetes	97.6 (97.3-98.0)	90.2 (89.5-90.8)	79.8 (78.9-80.7)	47.0 (45.8-48.3)	97.9 (97.6-98.1)	90.0 (89.4-90.7)	80.4 (79.5-81.2)
Hypertension / renal vascular disease	98.0 (97.7-98.4)	92.2 (91.4-92.9)	83.8 (82.8-84.9)	55.5 (53.8-57.2)	97.8 (97.4-98.2)	91.6 (90.8-92.4)	82.9 (81.8-84.0)
Glomerulonephritis	98.1 (97.7-98.6)	92.8 (91.9-93.7)	86.0 (84.6-87.3)	60.7 (58.5-63.1)	98.5 (98.1-98.9)	94.0 (93.2-94.8)	87.4 (86.2-88.6)
Other causes	96.7 (96.4-97.0)	89.1 (88.6-89.6)	80.1 (79.4-80.7)	52.6 (51.6-53.6)	97.0 (96.8-97.3)	89.4 (88.9-89.9)	80.6 (80.0-81.3)
All	97.3 (97.2-97.5)	90.3 (90.0-90.7)	81.4 (80.9-81.9)	52.7 (52.0-53.4)	97.6 (97.4-97.7)	90.5 (90.2-90.8)	81.8 (81.3-82.3)

Table B.6.7
Incident dialysis patients, from day 91, unadjusted
by age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	97.4 (96.4-98.1)	96.0 (94.6-97.0)	87.0 (82.8-90.3)	97.3 (96.3-98.1)	96.6 (95.3-97.5)
20-44	96.4 (96.0-96.7)	92.6 (92.1-93.1)	78.9 (77.9-79.9)	96.2 (95.9-96.5)	92.3 (91.8-92.7)
45-64	91.9 (91.6-92.1)	84.0 (83.6-84.3)	58.6 (58.0-59.1)	91.9 (91.6-92.1)	84.3 (83.9-84.6)
65-74	85.9 (85.5-86.2)	73.5 (73.1-74.0)	41.1 (40.5-41.6)	86.1 (85.8-86.4)	74.2 (73.7-74.6)
75+	78.6 (78.2-79.0)	61.5 (61.1-61.9)	25.0 (24.6-25.4)	79.3 (78.9-79.6)	62.3 (61.9-62.8)
Male	85.8 (85.6-86.0)	73.1 (72.8-73.4)	40.0 (39.6-40.3)	86.1 (85.8-86.3)	73.7 (73.4-74.0)
Female	86.7 (86.4-87.0)	75.2 (74.8-75.5)	44.3 (43.9-44.8)	87.2 (86.9-87.5)	75.9 (75.5-76.3)
Diabetes	86.6 (86.3-87.0)	73.3 (72.9-73.8)	38.0 (37.5-38.5)	86.8 (86.5-87.1)	73.6 (73.2-74.0)
Hypertension / renal vascular disease	85.3 (84.9-85.7)	72.2 (71.7-72.8)	37.7 (37.1-38.3)	85.6 (85.2-86.0)	72.8 (72.3-73.3)
Glomerulonephritis	92.9 (92.5-93.3)	85.5 (84.9-86.1)	59.0 (58.1-60.0)	93.1 (92.7-93.5)	86.1 (85.5-86.6)
Other causes	84.8 (84.5-85.1)	72.5 (72.1-72.8)	42.4 (41.9-42.8)	85.3 (85.0-85.5)	73.5 (73.1-73.8)
All	86.1 (86.0-86.3)	73.8 (73.6-74.1)	41.5 (41.2-41.8)	86.4 (86.3-86.6)	74.5 (74.3-74.7)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	97.9 (94.4-99.2)	96.4 (90.4-98.7)	93.3 (81.7-97.6)	98.3 (94.7-99.5)	98.3 (94.7-99.5)
20-44	97.5 (96.7-98.1)	95.4 (94.4-96.3)	83.7 (81.1-86.0)	97.7 (97.0-98.2)	95.7 (94.7-96.5)
45-64	93.6 (93.1-94.2)	86.9 (86.0-87.6)	60.4 (58.9-61.8)	93.7 (93.2-94.2)	87.3 (86.6-88.1)
65-74	88.9 (88.1-89.5)	77.1 (76.1-78.1)	44.6 (43.3-45.9)	87.9 (87.2-88.6)	77.0 (76.0-77.9)
75+	83.6 (82.8-84.4)	68.8 (67.8-69.8)	31.9 (30.9-33.0)	84.0 (83.2-84.7)	69.4 (68.4-70.3)
Male	89.1 (88.6-89.5)	77.4 (76.7-78.0)	43.0 (42.2-43.9)	88.7 (88.2-89.1)	77.7 (77.1-78.3)
Female	90.0 (89.4-90.6)	80.9 (80.0-81.7)	50.4 (49.1-51.6)	90.3 (89.7-90.9)	81.0 (80.2-81.9)
Diabetes	88.7 (87.9-89.4)	77.0 (75.9-78.0)	41.3 (39.9-42.6)	88.4 (87.7-89.1)	76.8 (75.8-77.7)
Hypertension / renal vascular disease	89.5 (88.4-90.4)	78.0 (76.5-79.3)	42.5 (40.7-44.3)	89.0 (87.9-90.0)	77.4 (76.0-78.8)
Glomerulonephritis	93.9 (93.0-94.7)	86.7 (85.3-87.9)	60.2 (57.7-62.5)	94.6 (93.7-95.3)	87.6 (86.3-88.7)
Other causes	88.7 (88.2-89.3)	77.9 (77.2-78.7)	46.6 (45.5-47.6)	88.6 (88.1-89.1)	78.6 (77.8-79.3)
All	89.4 (89.0-89.8)	78.5 (78.0-79.0)	45.4 (44.7-46.1)	89.2 (88.9-89.6)	78.8 (78.3-79.3)

Table B.6.8
Incident dialysis patients, from day 91, adjusted
adjusted for age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	97.3 (96.4-98.2)	95.7 (94.5-97.0)	86.4 (82.6-90.4)	97.2 (96.3-98.1)	96.3 (95.2-97.5)
20-44	96.3 (95.9-96.6)	92.4 (91.9-93.0)	78.6 (77.6-79.7)	96.2 (95.9-96.5)	92.2 (91.7-92.7)
45-64	92.1 (91.8-92.3)	84.3 (84.0-84.7)	59.2 (58.7-59.8)	92.1 (91.9-92.4)	84.7 (84.3-85.0)
65-74	86.4 (86.1-86.7)	74.4 (74.0-74.8)	42.2 (41.7-42.8)	86.6 (86.3-86.9)	75.0 (74.6-75.4)
75+	79.0 (78.7-79.4)	62.1 (61.7-62.5)	25.5 (25.1-25.9)	79.8 (79.5-80.2)	63.1 (62.7-63.6)
Male	87.7 (87.5-87.9)	76.4 (76.1-76.6)	44.8 (44.4-45.2)	88.0 (87.8-88.2)	77.1 (76.8-77.3)
Female	88.5 (88.3-88.8)	78.3 (78.0-78.7)	49.5 (49.0-50.0)	89.0 (88.7-89.2)	79.0 (78.7-79.4)
Diabetes	87.8 (87.5-88.1)	75.4 (75.0-75.8)	40.7 (40.1-41.2)	88.1 (87.8-88.4)	75.9 (75.5-76.3)
Hypertension / renal vascular disease	89.2 (88.9-89.6)	79.1 (78.7-79.5)	48.4 (47.8-49.1)	89.4 (89.1-89.7)	79.3 (78.9-79.7)
Glomerulonephritis	92.0 (91.6-92.5)	84.2 (83.6-84.8)	57.0 (56.0-58.0)	92.3 (91.9-92.7)	84.8 (84.2-85.4)
Other causes	86.4 (86.1-86.6)	75.2 (74.8-75.5)	46.4 (45.9-46.8)	86.9 (86.7-87.2)	76.3 (75.9-76.6)
All	88.0 (87.9-88.2)	77.1 (76.9-77.4)	46.6 (46.3-46.9)	88.4 (88.2-88.5)	77.8 (77.6-78.0)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	97.7 (95.6-100)	96.1 (92.5-100)	92.9 (86.0-100)	98.2 (96.2-100)	98.2 (96.2-100)
20-44	97.4 (96.8-98.1)	95.3 (94.4-96.3)	83.6 (81.1-86.1)	97.6 (96.9-98.2)	95.5 (94.6-96.5)
45-64	93.8 (93.3-94.3)	87.2 (86.4-88.0)	61.3 (59.8-62.8)	93.9 (93.4-94.4)	87.7 (86.9-88.4)
65-74	89.3 (88.6-90.0)	78.0 (77.1-79.0)	46.2 (44.9-47.5)	88.4 (87.7-89.0)	77.7 (76.8-78.7)
75+	83.9 (83.1-84.7)	69.4 (68.4-70.4)	32.7 (31.6-33.7)	84.5 (83.7-85.2)	70.2 (69.3-71.2)
Male	90.3 (89.9-90.7)	80.0 (79.4-80.6)	48.8 (47.9-49.7)	90.1 (89.7-90.6)	80.6 (80.0-81.2)
Female	91.2 (90.6-91.8)	83.3 (82.5-84.1)	56.8 (55.6-58.0)	91.6 (91.1-92.2)	83.7 (82.9-84.4)
Diabetes	89.8 (89.2-90.5)	79.2 (78.3-80.1)	45.6 (44.3-46.9)	89.9 (89.2-90.5)	79.6 (78.7-80.5)
Hypertension / renal vascular disease	92.1 (91.3-92.9)	83.4 (82.3-84.5)	53.7 (52.0-55.5)	91.8 (91.0-92.6)	83.0 (81.9-84.2)
Glomerulonephritis	93.2 (92.3-94.2)	85.6 (84.2-87.0)	59.7 (57.4-62.2)	94.0 (93.2-94.9)	86.7 (85.5-88.1)
Other causes	89.9 (89.4-90.4)	80.3 (79.6-81.0)	52.2 (51.1-53.2)	89.9 (89.4-90.4)	81.1 (80.4-81.8)
All	90.7 (90.3-91.0)	81.2 (80.8-81.7)	51.7 (51.0-52.5)	90.7 (90.4-91.1)	81.8 (81.3-82.2)

Table B.6.9

First transplant patients (deceased donor), from day of transplant, unadjusted

by age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	99.1 (98.3-99.5)	98.5 (97.6-99.1)	97.1 (95.9-98.0)	98.7 (97.8-99.2)	98.3 (97.4-98.9)
20-44	98.8 (98.6-99.0)	98.1 (97.8-98.4)	95.7 (95.3-96.1)	99.0 (98.7-99.2)	98.1 (97.8-98.4)
45-64	97.0 (96.8-97.3)	95.3 (95.0-95.6)	87.2 (86.8-87.7)	96.7 (96.5-97.0)	94.7 (94.4-95.0)
65-74	93.1 (92.6-93.6)	88.8 (88.1-89.4)	70.4 (69.5-71.4)	92.5 (91.9-93.0)	87.5 (86.8-88.2)
75+	90.7 (89.2-91.9)	83.4 (81.6-85.0)	57.5 (55.2-59.7)	89.1 (87.8-90.3)	80.3 (78.6-81.9)
Male	96.3 (96.1-96.6)	93.7 (93.4-94.0)	83.3 (82.9-83.8)	95.7 (95.5-96.0)	92.6 (92.3-93.0)
Female	96.3 (96.0-96.6)	94.6 (94.2-95.0)	86.2 (85.6-86.7)	96.1 (95.7-96.4)	93.7 (93.3-94.1)
Diabetes	94.5 (93.9-95.0)	90.8 (90.1-91.4)	75.7 (74.7-76.7)	93.3 (92.7-93.9)	88.7 (87.9-89.4)
Hypertension / renal vascular disease	95.6 (95.0-96.1)	92.0 (91.2-92.7)	79.1 (77.9-80.2)	94.9 (94.3-95.5)	90.9 (90.1-91.7)
Glomerulonephritis	97.4 (97.1-97.7)	95.9 (95.5-96.3)	88.7 (88.0-89.3)	97.3 (96.9-97.6)	95.4 (94.9-95.8)
Other causes	96.7 (96.5-97.0)	94.9 (94.6-95.2)	86.9 (86.4-87.3)	96.4 (96.2-96.7)	94.2 (93.9-94.5)
All	96.3 (96.1-96.5)	94.1 (93.8-94.3)	84.3 (84.0-84.7)	95.8 (95.6-96.0)	93.0 (92.8-93.3)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	99.5 (96.5-99.9)	99.5 (96.5-99.9)	98.0 (94.7-99.2)	99.6 (97.0-99.9)	99.6 (97.0-99.9)
20-44	98.5 (97.9-99.0)	98.2 (97.5-98.7)	96.0 (95.1-96.8)	99.1 (98.5-99.4)	98.5 (97.8-98.9)
45-64	96.8 (96.3-97.2)	95.3 (94.7-95.9)	87.6 (86.6-88.4)	96.5 (96.0-97.0)	94.7 (94.1-95.3)
65-74	91.7 (90.6-92.7)	88.0 (86.7-89.1)	69.0 (67.2-70.7)	92.0 (90.9-92.9)	87.4 (86.1-88.5)
75+	89.1 (86.3-91.3)	81.3 (78.0-84.2)	55.9 (51.9-59.7)	88.3 (85.9-90.3)	80.8 (78.0-83.3)
Male	95.5 (95.0-96.0)	93.1 (92.5-93.7)	81.9 (80.9-82.7)	95.0 (94.5-95.5)	92.1 (91.4-92.7)
Female	95.2 (94.4-95.8)	93.5 (92.7-94.2)	84.3 (83.1-85.4)	95.4 (94.7-96.1)	93.2 (92.4-94.0)
Diabetes	93.3 (92.1-94.3)	89.8 (88.3-91.1)	72.8 (70.7-74.8)	92.5 (91.2-93.5)	88.5 (87.0-89.8)
Hypertension / renal vascular disease	95.5 (94.1-96.6)	92.2 (90.4-93.6)	78.3 (75.8-80.7)	95.2 (93.8-96.4)	91.6 (89.8-93.1)
Glomerulonephritis	96.4 (95.5-97.1)	95.2 (94.2-96.1)	87.8 (86.3-89.1)	96.4 (95.6-97.1)	94.5 (93.5-95.4)
Other causes	95.7 (95.2-96.2)	93.9 (93.2-94.5)	84.9 (84.0-85.8)	95.7 (95.1-96.2)	93.3 (92.6-93.9)
All	95.4 (95.0-95.8)	93.2 (92.8-93.7)	82.7 (82.0-83.4)	95.2 (94.8-95.6)	92.5 (91.9-92.9)

Table B.6.10

First transplant patients (deceased donor), from day of transplant, adjusted

adjusted for age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	99.0 (98.3-99.6)	98.4 (97.6-99.2)	96.8 (95.7-97.9)	98.6 (97.8-99.3)	98.1 (97.3-99.0)
20-44	98.8 (98.6-99.1)	98.2 (97.9-98.5)	95.8 (95.4-96.2)	99.0 (98.8-99.2)	98.2 (97.9-98.4)
45-64	97.2 (97.0-97.5)	95.6 (95.4-95.9)	88.0 (87.6-88.5)	97.0 (96.7-97.2)	95.1 (94.8-95.4)
65-74	93.8 (93.3-94.2)	89.8 (89.2-90.4)	72.7 (71.7-73.6)	93.2 (92.7-93.7)	88.7 (88.0-89.3)
75+	91.5 (90.3-92.7)	84.8 (83.3-86.4)	60.2 (58.0-62.4)	90.3 (89.1-91.4)	82.3 (80.8-83.8)
Male	98.2 (98.1-98.4)	97.0 (96.8-97.1)	91.3 (91.0-91.6)	98.1 (98.0-98.2)	96.7 (96.5-96.8)
Female	98.0 (97.9-98.2)	97.1 (96.9-97.3)	92.1 (91.7-92.5)	98.0 (97.8-98.2)	96.8 (96.5-97.0)
Diabetes	97.3 (97.0-97.5)	95.3 (95.0-95.7)	86.7 (86.0-87.4)	96.9 (96.6-97.2)	94.7 (94.3-95.1)
Hypertension / renal vascular disease	98.3 (98.0-98.5)	96.8 (96.4-97.1)	90.8 (90.2-91.4)	98.1 (97.8-98.3)	96.5 (96.1-96.8)
Glomerulonephritis	98.4 (98.2-98.7)	97.5 (97.2-97.8)	92.8 (92.3-93.3)	98.4 (98.2-98.6)	97.3 (97.1-97.6)
Other causes	98.1 (98.0-98.3)	97.1 (96.9-97.3)	92.1 (91.7-92.4)	98.1 (97.9-98.2)	96.9 (96.6-97.1)
All	98.2 (98.1-98.3)	97.0 (96.9-97.2)	91.6 (91.3-91.9)	98.1 (98.0-98.2)	96.7 (96.5-96.9)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	99.5 (98.4-100)	99.5 (98.4-100)	97.8 (95.7-100)	99.5 (98.6-100)	99.5 (98.6-100)
20-44	98.6 (98.0-99.1)	98.2 (97.6-98.8)	96.1 (95.3-97.0)	99.1 (98.6-99.5)	98.5 (97.9-99.0)
45-64	97.0 (96.5-97.4)	95.6 (95.1-96.1)	88.2 (87.3-89.1)	96.7 (96.2-97.1)	95.0 (94.4-95.6)
65-74	92.4 (91.4-93.4)	89.0 (87.9-90.1)	71.0 (69.4-72.7)	92.7 (91.8-93.6)	88.5 (87.4-89.6)
75+	90.0 (87.8-92.3)	82.8 (80.0-85.7)	58.4 (54.7-62.3)	89.1 (87.1-91.2)	82.2 (79.7-84.8)
Male	98.2 (98.0-98.4)	97.2 (96.9-97.5)	91.9 (91.3-92.5)	98.1 (97.8-98.3)	96.9 (96.5-97.2)
Female	97.9 (97.6-98.2)	97.2 (96.8-97.6)	92.7 (92.0-93.4)	98.1 (97.7-98.4)	97.1 (96.7-97.5)
Diabetes	97.4 (96.9-97.9)	95.9 (95.3-96.6)	87.9 (86.7-89.1)	97.2 (96.7-97.7)	95.6 (95.0-96.3)
Hypertension / renal vascular disease	98.6 (98.2-99.0)	97.5 (96.9-98.0)	92.3 (91.2-93.4)	98.5 (98.0-98.9)	97.3 (96.7-97.9)
Glomerulonephritis	98.1 (97.7-98.5)	97.5 (97.0-98.0)	93.2 (92.3-94.1)	98.2 (97.8-98.6)	97.2 (96.7-97.8)
Other causes	98.1 (97.8-98.3)	97.2 (96.9-97.5)	92.6 (92.0-93.3)	98.1 (97.8-98.4)	97.0 (96.7-97.4)
All	98.1 (97.9-98.3)	97.2 (97.0-97.5)	92.2 (91.7-92.8)	98.1 (97.9-98.3)	97.0 (96.7-97.3)

Table B.6.11
First transplant patients (living donor), from day of transplant, unadjusted
by age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	99.2 (98.4-99.6)	99.0 (98.1-99.5)	98.3 (97.2-98.9)	99.4 (98.6-99.8)	99.0 (98.1-99.5)
20-44	99.6 (99.4-99.7)	99.3 (99.0-99.5)	98.4 (98.0-98.7)	99.7 (99.5-99.9)	99.4 (99.0-99.6)
45-64	98.8 (98.5-99.0)	97.9 (97.4-98.2)	93.3 (92.6-94.0)	98.7 (98.3-99.0)	97.6 (97.1-98.0)
65-74	97.0 (96.1-97.8)	94.6 (93.3-95.6)	83.4 (81.4-85.2)	97.2 (96.2-97.9)	95.1 (93.9-96.1)
75+	94.6 (89.0-97.4)	93.0 (87.0-96.3)	68.0 (59.1-75.3)	93.9 (89.0-96.7)	90.9 (85.4-94.4)
Male	98.6 (98.4-98.9)	97.8 (97.5-98.1)	93.8 (93.2-94.3)	98.6 (98.3-98.8)	97.6 (97.3-98.0)
Female	99.2 (98.9-99.4)	98.3 (97.9-98.6)	94.5 (93.7-95.1)	99.3 (98.9-99.5)	98.3 (97.9-98.7)
Diabetes	97.3 (96.1-98.2)	94.8 (93.3-96.0)	84.2 (81.8-86.4)	97.9 (96.7-98.7)	95.5 (93.9-96.7)
Hypertension / renal vascular disease	97.9 (96.9-98.7)	96.2 (94.8-97.2)	89.4 (87.3-91.2)	97.1 (95.8-98.0)	95.8 (94.3-96.9)
Glomerulonephritis	99.0 (98.6-99.3)	98.6 (98.1-99.0)	96.2 (95.4-96.8)	99.3 (98.9-99.6)	98.8 (98.3-99.1)
Other causes	99.1 (98.9-99.3)	98.5 (98.2-98.8)	95.2 (94.7-95.7)	99.0 (98.7-99.2)	98.1 (97.8-98.4)
All	98.8 (98.6-99.0)	98.0 (97.8-98.2)	94.0 (93.6-94.4)	98.8 (98.6-99.0)	97.9 (97.6-98.1)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	100 (100.0-100)	100 (100.0-100)	99.4 (98.3-99.8)	99.8 (98.4-100)	99.8 (98.4-100)
45-64	99.2 (98.1-99.7)	98.1 (96.7-98.9)	94.3 (92.2-95.9)	99.5 (98.3-99.8)	98.6 (97.1-99.3)
65-74	96.7 (92.2-98.6)	93.4 (88.1-96.4)	86.1 (79.4-90.7)	96.1 (91.6-98.2)	92.9 (87.6-96.0)
75+					
Male	99.0 (98.1-99.5)	98.1 (97.0-98.8)	95.3 (93.7-96.5)	98.9 (97.9-99.4)	97.8 (96.5-98.6)
Female	99.8 (98.7-100)	99.1 (97.8-99.6)	96.4 (94.5-97.7)	99.8 (98.4-100)	99.3 (97.9-99.8)
Diabetes	100 (100.0-100)	96.4 (90.8-98.6)	91.9 (85.1-95.7)	100 (100.0-100)	97.0 (91.0-99.0)
Hypertension / renal vascular disease					
Glomerulonephritis	98.9 (97.0-99.6)	98.3 (96.3-99.2)	96.9 (94.5-98.3)	98.7 (96.5-99.5)	98.0 (95.6-99.1)
Other causes	99.7 (99.0-99.9)	99.2 (98.4-99.6)	96.3 (94.8-97.3)	99.3 (98.4-99.7)	99.0 (97.9-99.5)
All	99.3 (98.7-99.6)	98.5 (97.7-99.0)	95.7 (94.6-96.7)	99.2 (98.5-99.6)	98.3 (97.4-98.9)

Table B.6.12
First transplant patients (living donor), from day of transplant, adjusted
adjusted for age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	99.2 (98.6-99.8)	99.0 (98.2-99.7)	98.0 (97.0-99.0)	99.3 (98.7-99.9)	98.9 (98.2-99.7)
20-44	99.6 (99.4-99.8)	99.3 (99.0-99.5)	98.3 (97.9-98.7)	99.7 (99.5-99.9)	99.3 (99.0-99.6)
45-64	98.8 (98.5-99.1)	97.9 (97.6-98.3)	93.6 (92.9-94.2)	98.7 (98.4-99.0)	97.6 (97.2-98.1)
65-74	97.4 (96.6-98.1)	95.1 (94.1-96.2)	84.9 (83.2-86.7)	97.5 (96.8-98.3)	95.7 (94.7-96.7)
75+	95.2 (91.8-98.7)	93.9 (90.0-97.8)	70.6 (63.4-78.7)	94.5 (91.3-97.9)	91.8 (87.8-95.9)
Male	99.0 (98.8-99.2)	98.4 (98.2-98.7)	95.3 (94.8-95.7)	98.9 (98.7-99.2)	98.2 (97.9-98.5)
Female	99.3 (99.1-99.6)	98.6 (98.2-98.9)	95.1 (94.5-95.8)	99.4 (99.1-99.6)	98.6 (98.3-98.9)
Diabetes	98.4 (97.8-99.0)	96.9 (96.1-97.7)	90.1 (88.5-91.6)	98.8 (98.3-99.4)	97.4 (96.6-98.3)
Hypertension / renal vascular disease	98.9 (98.4-99.4)	97.9 (97.2-98.6)	94.0 (92.8-95.2)	98.5 (97.9-99.1)	97.8 (97.1-98.5)
Glomerulonephritis	99.1 (98.7-99.4)	98.6 (98.2-99.0)	96.1 (95.4-96.8)	99.3 (99.1-99.6)	98.8 (98.4-99.2)
Other causes	99.2 (99.1-99.4)	98.7 (98.5-99.0)	95.8 (95.3-96.3)	99.1 (98.9-99.3)	98.4 (98.1-98.7)
All	99.1 (99.0-99.3)	98.5 (98.3-98.7)	95.2 (94.8-95.6)	99.1 (98.9-99.3)	98.4 (98.1-98.6)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	100	100	99.4 (98.7-100)	99.7 (99.3-100)	99.7 (99.3-100)
45-64	99.2 (98.5-99.9)	98.0 (97.0-99.1)	94.3 (92.5-96.1)	99.5 (98.8-100)	98.5 (97.5-99.6)
65-74	96.8 (94.1-99.6)	93.6 (89.9-97.6)	86.6 (81.3-92.2)	96.5 (93.8-99.3)	93.6 (89.9-97.4)
75+					
Male	99.4 (99.0-99.8)	98.8 (98.2-99.4)	97.1 (96.1-98.1)	99.5 (99.2-99.9)	99.1 (98.5-99.6)
Female	99.8 (99.5-100)	99.2 (98.4-99.9)	96.7 (95.2-98.2)	99.9 (99.6-100)	99.6 (99.1-100)
Diabetes	100	98.4 (96.8-100)	96.3 (93.8-98.8)	100	99.2 (98.3-100)
Hypertension / renal vascular disease					
Glomerulonephritis	99.0 (98.0-100)	98.5 (97.3-99.7)	97.2 (95.6-98.9)	99.2 (98.3-100)	98.7 (97.7-99.8)
Other causes	99.7 (99.5-100)	99.4 (99.0-99.9)	97.1 (96.0-98.2)	99.6 (99.3-100)	99.4 (98.9-99.9)
All	99.5 (99.2-99.8)	98.9 (98.4-99.4)	97.0 (96.1-97.9)	99.6 (99.3-99.9)	99.2 (98.7-99.7)

Table B.6.13
First graft (deceased donor), from day of transplant, unadjusted
by age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	94.8 (93.2-95.9)	92.3 (90.6-93.8)	84.9 (82.6-86.9)	94.4 (92.8-95.6)	91.8 (90.0-93.3)
20-44	94.5 (94.0-94.9)	92.5 (92.0-93.0)	85.2 (84.5-85.9)	95.3 (94.9-95.7)	93.2 (92.7-93.7)
45-64	92.2 (91.8-92.5)	89.3 (88.9-89.8)	78.8 (78.3-79.4)	92.1 (91.7-92.5)	89.3 (88.8-89.7)
65-74	86.5 (85.7-87.2)	81.3 (80.5-82.1)	62.0 (61.0-63.0)	86.4 (85.7-87.0)	80.9 (80.1-81.7)
75+	83.1 (81.3-84.7)	75.1 (73.1-77.0)	50.1 (47.8-52.4)	81.7 (80.0-83.2)	72.2 (70.3-73.9)
Male	90.9 (90.6-91.3)	87.4 (86.9-87.7)	74.5 (73.9-75.0)	90.6 (90.2-90.9)	86.7 (86.3-87.1)
Female	91.4 (90.9-91.8)	88.4 (87.9-88.9)	77.1 (76.4-77.7)	91.6 (91.1-92.0)	88.2 (87.6-88.7)
Diabetes	89.5 (88.8-90.2)	85.1 (84.3-85.9)	68.6 (67.5-69.7)	88.1 (87.4-88.9)	82.9 (82.0-83.7)
Hypertension / renal vascular disease	89.4 (88.5-90.2)	84.7 (83.6-85.6)	69.0 (67.7-70.3)	89.6 (88.7-90.4)	84.6 (83.6-85.6)
Glomerulonephritis	91.8 (91.2-92.3)	89.0 (88.3-89.6)	78.3 (77.4-79.2)	92.1 (91.5-92.7)	89.2 (88.6-89.9)
Other causes	91.8 (91.4-92.1)	88.9 (88.5-89.3)	78.2 (77.6-78.7)	91.8 (91.4-92.2)	88.7 (88.3-89.1)
All	91.1 (90.8-91.4)	87.7 (87.4-88.0)	75.4 (75.0-75.8)	90.9 (90.6-91.2)	87.2 (86.9-87.6)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	93.0 (88.5-95.8)	90.0 (84.9-93.4)	81.3 (75.1-86.1)	93.2 (89.1-95.8)	90.2 (85.6-93.4)
20-44	92.5 (91.2-93.5)	90.9 (89.5-92.1)	84.0 (82.3-85.5)	94.1 (92.9-95.1)	91.9 (90.5-93.1)
45-64	90.8 (90.0-91.6)	88.2 (87.3-89.0)	77.8 (76.6-78.9)	90.4 (89.6-91.1)	87.6 (86.7-88.5)
65-74	83.4 (82.0-84.8)	78.8 (77.2-80.3)	59.3 (57.5-61.1)	84.4 (83.0-85.7)	79.4 (77.9-80.8)
75+	79.9 (76.5-82.8)	71.7 (68.0-75.1)	48.2 (44.2-52.0)	80.4 (77.5-82.9)	71.9 (68.7-74.8)
Male	88.7 (87.9-89.4)	85.1 (84.2-85.9)	71.5 (70.4-72.5)	88.3 (87.5-89.0)	84.4 (83.6-85.3)
Female	88.5 (87.5-89.5)	85.9 (84.7-86.9)	74.5 (73.1-75.9)	89.6 (88.6-90.5)	86.1 (85.0-87.2)
Diabetes	86.8 (85.2-88.3)	82.9 (81.2-84.6)	65.0 (62.7-67.1)	85.7 (84.1-87.1)	81.5 (79.7-83.1)
Hypertension / renal vascular disease	87.2 (85.0-89.0)	82.5 (80.1-84.6)	66.8 (63.9-69.5)	88.2 (86.2-90.0)	83.3 (80.9-85.4)
Glomerulonephritis	89.7 (88.3-91.0)	87.4 (85.9-88.8)	76.3 (74.4-78.0)	89.7 (88.4-90.9)	86.8 (85.3-88.1)
Other causes	89.1 (88.3-89.9)	85.9 (85.0-86.8)	74.7 (73.6-75.8)	89.6 (88.7-90.3)	86.0 (85.1-86.9)
All	88.6 (88.0-89.2)	85.3 (84.7-86.0)	72.5 (71.7-73.3)	88.7 (88.1-89.3)	85.0 (84.3-85.7)

Table B.6.14

First graft (deceased donor), from day of transplant, adjusted

adjusted for age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	94.3 (92.9-95.7)	91.7 (90.0-93.4)	83.6 (81.4-86.0)	93.9 (92.4-95.4)	91.1 (89.4-92.9)
20-44	94.5 (94.1-95.0)	92.6 (92.1-93.1)	85.3 (84.6-86.1)	95.4 (94.9-95.8)	93.3 (92.7-93.8)
45-64	92.4 (92.1-92.8)	89.7 (89.3-90.1)	79.5 (78.9-80.0)	92.4 (92.0-92.8)	89.7 (89.2-90.1)
65-74	87.2 (86.5-87.9)	82.3 (81.6-83.1)	63.6 (62.6-64.6)	87.1 (86.5-87.8)	81.9 (81.1-82.6)
75+	84.0 (82.4-85.7)	76.4 (74.5-78.3)	52.0 (49.7-54.3)	82.7 (81.3-84.3)	73.7 (71.9-75.5)
Male	93.3 (93.0-93.6)	90.6 (90.2-90.9)	80.2 (79.7-80.7)	93.3 (93.0-93.5)	90.4 (90.1-90.8)
Female	93.2 (92.8-93.6)	90.8 (90.3-91.2)	81.2 (80.6-81.8)	93.5 (93.2-93.9)	90.8 (90.4-91.3)
Diabetes	92.2 (91.7-92.8)	88.8 (88.2-89.5)	75.3 (74.4-76.3)	91.6 (91.0-92.1)	87.6 (86.9-88.3)
Hypertension / renal vascular disease	93.0 (92.4-93.6)	89.7 (89.0-90.4)	78.2 (77.2-79.2)	93.3 (92.7-93.9)	90.0 (89.2-90.7)
Glomerulonephritis	93.1 (92.6-93.6)	90.7 (90.1-91.3)	81.3 (80.5-82.1)	93.6 (93.1-94.0)	91.2 (90.6-91.7)
Other causes	93.3 (93.0-93.6)	90.9 (90.5-91.3)	81.7 (81.1-82.2)	93.5 (93.2-93.8)	91.0 (90.6-91.4)
All	93.3 (93.0-93.5)	90.6 (90.4-90.9)	80.6 (80.2-81.0)	93.4 (93.1-93.6)	90.6 (90.3-90.9)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	92.6 (88.9-96.4)	89.4 (85.1-93.9)	80.3 (74.7-86.2)	92.9 (89.5-96.3)	89.7 (85.8-93.8)
20-44	92.5 (91.3-93.6)	90.9 (89.6-92.2)	84.0 (82.4-85.6)	94.2 (93.1-95.2)	91.9 (90.7-93.2)
45-64	91.1 (90.3-91.8)	88.5 (87.6-89.3)	78.2 (77.1-79.4)	90.6 (89.8-91.4)	87.9 (87.0-88.8)
65-74	84.1 (82.8-85.5)	79.7 (78.2-81.2)	60.6 (58.8-62.5)	85.1 (83.8-86.4)	80.2 (78.8-81.7)
75+	80.8 (77.8-83.9)	72.8 (69.4-76.4)	49.5 (45.8-53.6)	81.0 (78.4-83.7)	72.7 (69.8-75.8)
Male	92.1 (91.5-92.7)	89.5 (88.8-90.1)	78.9 (77.9-79.9)	92.0 (91.4-92.6)	89.2 (88.5-89.9)
Female	91.6 (90.8-92.4)	89.6 (88.8-90.5)	80.6 (79.4-81.9)	92.4 (91.6-93.2)	89.8 (88.9-90.7)
Diabetes	91.1 (90.0-92.2)	88.4 (87.1-89.6)	74.5 (72.7-76.4)	90.6 (89.5-91.7)	87.7 (86.4-89.0)
Hypertension / renal vascular disease	92.2 (90.9-93.5)	89.2 (87.7-90.7)	78.3 (76.2-80.4)	92.8 (91.6-94.1)	89.6 (88.1-91.2)
Glomerulonephritis	91.8 (90.8-92.9)	90.0 (88.8-91.2)	80.6 (79.0-82.2)	92.0 (91.0-93.0)	89.6 (88.5-90.8)
Other causes	91.9 (91.3-92.6)	89.5 (88.7-90.2)	80.4 (79.4-81.5)	92.3 (91.7-93.0)	89.6 (88.9-90.4)
All	92.0 (91.5-92.5)	89.6 (89.0-90.1)	79.6 (78.7-80.4)	92.1 (91.6-92.7)	89.5 (88.8-90.1)

Table B.6.15
First graft (living donor), from day of transplant, unadjusted
by age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	96.0 (94.5-97.1)	94.2 (92.4-95.5)	89.3 (87.1-91.2)	97.5 (96.2-98.4)	95.7 (94.1-96.9)
20-44	96.8 (96.3-97.3)	95.7 (95.1-96.3)	90.0 (89.0-90.8)	97.6 (97.0-98.0)	96.3 (95.6-96.8)
45-64	96.7 (96.2-97.1)	95.2 (94.6-95.7)	88.8 (88.0-89.7)	96.8 (96.3-97.3)	95.2 (94.6-95.8)
65-74	95.5 (94.3-96.4)	92.7 (91.3-93.9)	80.2 (78.1-82.1)	95.8 (94.7-96.7)	93.2 (91.8-94.3)
75+	94.6 (89.0-97.4)	92.2 (86.1-95.8)	66.4 (57.5-73.9)	93.3 (88.3-96.3)	90.3 (84.7-93.9)
Male	96.4 (96.0-96.8)	95.0 (94.5-95.5)	88.3 (87.6-89.0)	96.9 (96.4-97.2)	95.3 (94.8-95.8)
Female	96.7 (96.2-97.2)	94.9 (94.2-95.5)	87.4 (86.4-88.4)	97.1 (96.5-97.6)	95.2 (94.5-95.8)
Diabetes	94.9 (93.3-96.1)	91.5 (89.6-93.0)	79.6 (76.9-81.9)	96.1 (94.6-97.2)	93.0 (91.1-94.5)
Hypertension / renal vascular disease	96.0 (94.6-97.0)	93.1 (91.4-94.5)	82.9 (80.4-85.1)	95.4 (93.8-96.5)	92.9 (91.0-94.4)
Glomerulonephritis	96.6 (95.9-97.2)	95.5 (94.7-96.2)	88.5 (87.3-89.6)	97.2 (96.5-97.7)	95.8 (95.0-96.5)
Other causes	96.8 (96.4-97.2)	95.5 (95.0-96.0)	89.8 (89.0-90.4)	97.1 (96.7-97.5)	95.7 (95.1-96.1)
All	96.5 (96.2-96.8)	95.0 (94.6-95.3)	88.0 (87.4-88.5)	96.9 (96.6-97.2)	95.3 (94.9-95.7)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	95.5 (93.4-96.9)	94.8 (92.6-96.3)	90.2 (87.4-92.4)	97.3 (95.3-98.5)	96.2 (93.9-97.6)
45-64	96.0 (94.2-97.3)	94.5 (92.4-96.0)	89.2 (86.6-91.4)	95.5 (93.4-96.9)	94.2 (91.9-95.9)
65-74	93.4 (88.1-96.4)	90.1 (84.1-93.9)	83.4 (76.5-88.5)	94.2 (89.2-97.0)	91.0 (85.3-94.6)
75+					
Male	95.1 (93.5-96.3)	93.6 (91.8-95.0)	88.9 (86.6-90.8)	96.3 (94.7-97.4)	94.8 (93.0-96.1)
Female	96.7 (94.7-97.9)	95.0 (92.8-96.5)	89.4 (86.4-91.7)	96.2 (94.0-97.6)	94.4 (91.9-96.2)
Diabetes	94.6 (88.5-97.6)	91.1 (84.0-95.1)	86.6 (78.7-91.7)	100 (100.0-100)	97.0 (91.0-99.0)
Hypertension / renal vascular disease					
Glomerulonephritis	96.1 (93.5-97.7)	95.2 (92.5-97.0)	90.5 (86.9-93.1)	96.0 (93.1-97.7)	95.0 (91.9-97.0)
Other causes	95.9 (94.4-97.0)	94.3 (92.6-95.6)	89.1 (86.9-91.0)	96.0 (94.3-97.1)	94.5 (92.6-95.9)
All	95.7 (94.5-96.6)	94.1 (92.7-95.2)	89.1 (87.3-90.6)	96.2 (95.0-97.2)	94.6 (93.2-95.8)

Table B.6.16
First graft (living donor), from day of transplant, adjusted
adjusted for age, sex, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19	95.4 (94.0-96.9)	93.2 (91.5-95.0)	87.7 (85.5-90.1)	97.1 (95.9-98.3)	95.0 (93.5-96.6)
20-44	96.7 (96.2-97.3)	95.6 (95.0-96.2)	89.5 (88.6-90.4)	97.4 (96.9-97.9)	96.0 (95.4-96.7)
45-64	96.7 (96.2-97.2)	95.2 (94.6-95.8)	88.9 (88.1-89.7)	96.8 (96.3-97.3)	95.2 (94.5-95.8)
65-74	95.8 (94.8-96.8)	93.1 (91.8-94.4)	81.1 (79.1-83.1)	96.1 (95.1-97.0)	93.6 (92.3-94.8)
75+	94.9 (91.3-98.7)	92.8 (88.5-97.2)	67.7 (60.1-76.2)	93.5 (89.8-97.3)	90.5 (86.2-95.1)
Male	96.4 (96.0-96.8)	95.0 (94.5-95.5)	88.1 (87.4-88.8)	96.8 (96.4-97.2)	95.3 (94.8-95.8)
Female	96.5 (96.0-97.1)	94.5 (93.8-95.2)	86.6 (85.5-87.6)	96.9 (96.3-97.4)	94.8 (94.1-95.5)
Diabetes	95.3 (94.0-96.5)	92.1 (90.5-93.7)	80.9 (78.6-83.3)	96.5 (95.3-97.7)	93.7 (92.1-95.2)
Hypertension / renal vascular disease	96.3 (95.2-97.4)	93.7 (92.3-95.1)	84.1 (82.0-86.4)	95.9 (94.7-97.1)	93.7 (92.2-95.2)
Glomerulonephritis	96.4 (95.7-97.0)	95.1 (94.3-95.9)	87.6 (86.4-88.8)	97.0 (96.3-97.7)	95.5 (94.7-96.3)
Other causes	96.7 (96.3-97.1)	95.4 (94.9-95.9)	89.4 (88.6-90.1)	97.0 (96.6-97.4)	95.5 (95.0-96.0)
All	96.4 (96.1-96.8)	94.8 (94.4-95.2)	87.6 (87.0-88.2)	96.8 (96.5-97.2)	95.1 (94.7-95.5)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Bosnia and Herzegovina, Denmark, Estonia, France, Greece, Iceland, Netherlands, Norway, Spain (Andalusia), Spain (Aragon), Spain (Basque country), Spain (Canary Islands), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Community of Madrid), Spain (Extremadura), Spain (Galicia), Spain (Murcia), Spain (Navarre), Spain (Valencian Region), Sweden, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Spain	Survival probabilities as % (95% CI)				
	cohort 2014 - 2018			cohort 2017 - 2021	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	95.5 (93.7-97.3)	94.7 (92.8-96.6)	89.9 (87.3-92.6)	97.3 (95.7-98.8)	96.1 (94.3-98.0)
45-64	96.0 (94.4-97.5)	94.3 (92.5-96.2)	89.2 (86.7-91.7)	95.4 (93.7-97.2)	94.2 (92.2-96.1)
65-74	93.4 (89.6-97.5)	90.1 (85.5-95.0)	83.5 (77.8-89.7)	94.3 (90.7-98.0)	91.0 (86.6-95.7)
75+					
Male	95.1 (93.7-96.6)	93.5 (91.9-95.2)	88.8 (86.7-90.9)	96.5 (95.2-97.7)	95.0 (93.6-96.6)
Female	96.3 (94.6-98.0)	94.4 (92.4-96.5)	88.1 (85.3-91.1)	96.0 (94.1-97.9)	94.1 (91.8-96.4)
Diabetes	95.3 (91.7-99.1)	92.2 (87.6-97.0)	88.2 (82.7-94.0)	100	97.8 (95.4-100)
Hypertension / renal vascular disease					
Glomerulonephritis	95.8 (93.6-98.0)	94.9 (92.5-97.3)	89.7 (86.5-93.1)	95.8 (93.6-98.2)	94.8 (92.3-97.4)
Other causes	95.7 (94.3-97.0)	94.0 (92.4-95.6)	88.6 (86.5-90.7)	95.8 (94.4-97.3)	94.4 (92.7-96.0)
All	95.5 (94.4-96.6)	93.8 (92.5-95.1)	88.6 (86.9-90.3)	96.3 (95.2-97.4)	94.7 (93.5-96.0)